

The multiple faces of caveolae

Robert G. Parton* and Kai Simons†

Abstract | Caveolae are a highly abundant but enigmatic feature of mammalian cells. They form remarkably stable membrane domains at the plasma membrane but can also function as carriers in the exocytic and endocytic pathways. The apparently diverse functions of caveolae, including mechanosensing and lipid regulation, might be linked to their ability to respond to plasma membrane changes, a property that is dependent on their specialized lipid composition and biophysical properties.

Stomatal diaphragm

A specialized structure at the neck of caveolae in certain endothelial cells that consists of a central density and radial spikes, and is generated by the transmembrane protein PV1.

Lipid rafts

Small, heterogeneous, highly dynamic, sterol- and sphingolipid-enriched domains that are formed by lipid–lipid interactions that compartmentalize cellular processes. Small lipid rafts can be stabilized to form larger platforms through protein–protein and protein–lipid interactions.

Caveolae were first identified by electron microscopy^{1,2} and are defined as pits of 60–80-nm diameter in the plasma membrane (FIG. 1). Caveolae have a characteristic flask shape and no obvious coat³, as is seen, for example, on the cytoplasmic surface of clathrin-coated pits. In some tissues, caveolae have a unique specialized structure at the neck of the flask, known as the stomatal diaphragm⁴.

The incredible abundance of caveolae in specific cell types, such as smooth-muscle cells, fibroblasts (BOX 1; FIG. 1), endothelial cells and adipocytes has long puzzled researchers who are seeking to understand their cellular functions. Caveolae have been implicated in endocytosis, transcytosis, calcium signalling and numerous other signal transduction events^{5–7}. They have also been exploited by pathogens, both as direct portals for endocytic entry⁸ and to facilitate the entry of pathogens that are larger than a single caveola⁹. Caveolae, and the main membrane proteins of caveolae, caveolins, have also been linked to disease; mutations in caveolins have been found in breast cancer and in limb girdle muscular dystrophy (BOX 2).

What might the function of this specialized surface domain be? In the past, signalling has been proposed as the primary function of caveolae. Although caveolae are certainly involved in specific signalling events, as will be discussed below, a more general role in signalling remains controversial and must be reinvestigated with the functional systems that are now available. The ability to experimentally manipulate caveolins, which are essential for caveola formation^{10–12}, has allowed researchers to examine the effect of caveola deficiency in specific cell types in tissue culture and *in vivo*. These studies are providing fascinating new insights into the functions of caveolae.

Features and properties of caveolins

Caveolin-1 (CAV1) and CAV2 are abundant in caveola-rich non-muscle cells, whereas CAV3 is found in skeletal muscle and in some smooth-muscle cells^{13,14}. Ablation

of CAV1 and CAV3 causes loss of caveolae from those specific cell types^{11,12}. By contrast, loss of CAV2 has no apparent effect on caveola formation *in vivo*¹⁵ but might contribute to caveola formation in certain cell types^{16,17}. All three caveolins show an unusual topology with N and C termini in the cytoplasm and a long putative hairpin intramembrane domain. Caveola formation by CAV1 and CAV3 involves oligomerization and association with cholesterol-rich lipid-raft domains. CAV1 binds to 1–2 cholesterol molecules¹⁸ and is also palmitoylated in the C-terminal region¹⁹. Cholesterol depletion disrupts the structure of caveolae²⁰.

The main structural features of caveolin are summarized in FIG. 1. An estimated 144 molecules of caveolin are present in a single caveolar structure²¹. The relative amount of cholesterol in caveolae might be more than 100 times higher than this, estimated as 20,000 molecules in immuno-isolated caveolae²². Some glycosphingolipids (for example, GM1 and GM3) and sphingomyelin are also enriched in caveolae relative to the bulk plasma membrane. Importantly, the density of lipids was found to be higher in the caveolae than in the plasma membrane fraction from which the caveolae were isolated²². Both the lipid composition and the packing of the lipid bilayer conform to the parameters that would be expected from a clustered liquid-ordered lipid-raft domain^{23–25}. In summary, caveolae represent a specialized, morphologically distinct sphingolipid-cholesterol microdomain, which is stabilized by the caveolin protein.

Trafficking of caveolins

The remarkable ability of exogeneously expressed CAV1 or CAV3 to generate caveolae has been shown in many experimental systems^{10,26} (for a review addressing mechanisms of caveola formation, see REF. 27). An understanding of caveolin trafficking and caveola formation is crucial to understanding the possible role of caveolins and caveolae.

*Institute for Molecular Bioscience and Centre for Microscopy and Microanalysis, University of Queensland, Brisbane, Queensland 4072, Australia.

†Max Planck Institute for Molecular Cell Biology and Genetics, Pflotenhauerstrasse 110, D-01307 Dresden, Germany.
e-mails:
r.parton@imb.uq.edu.au;
simons@mpi-cbg.de
doi:10.1038/nrm2122




Figure 1 | Caveolae and caveolins. The diagrams show the main features of caveolae and caveolins. The electron micrographs in panels **a** and **b** show caveolae in adipocytes that have been surface-labelled with an electron-dense marker. Panel **c** shows a glancing section across the cell surface of a primary fibroblast that has been similarly labelled. Caveolae are evident as discrete flask-shaped pits, or circular profiles where the surface connection lies outside the plane of the section. Note the complex forms of surface-connected caveolae in the adipocytes (panels **a** and **b**), and the incredible abundance of caveolae in specific regions of the fibroblast surface (panel **c**). Panel **d** indicates how caveolin is inserted into the caveolar membrane, with the N and C termini facing the cytoplasm and a putative 'hairpin' intramembrane domain embedded within the membrane bilayer. The scaffolding domain, a highly conserved region of caveolin, might have a role in cholesterol interactions through conserved basic (+) and bulky hydrophobic residues (red circles). The C-terminal domain, which is close to the intramembrane domain, is modified by palmitoyl groups that insert into the lipid bilayer. The complex structures that are formed by interconnected caveolae can occupy a large area of the plasma membrane. The hypothetical formation of cubic membranes (panel **e**), which have adapted to allow the invagination of numerous caveolae, is depicted schematically¹⁰⁸. These membrane invaginations can form with little energy input. Panel **e** is reproduced with permission from REF. 108 © (1995) Elsevier.

Synthesis and trafficking of caveolins. Caveolin is synthesized as an integral membrane protein in the endoplasmic reticulum (ER) in a signal recognition particle (SRP)-dependent manner²⁸. The newly synthesized protein goes through a first stage of oligomerization which, at least *in vitro*, can occur in the ER²⁹. Caveolin is then transported from the ER to the Golgi complex. A Golgi pool of newly synthesized caveolin is observed in many cell types, and this pool of the protein is not associated with detergent-resistant membrane (DRM)³⁰. However, at some point in the biosynthetic pathway, caveolin associates with lipid rafts, becomes detergent-resistant and is organized into higher-order oligomers that are characteristic of the surface pool of the protein.

The fact that the Golgi and plasma membrane pools of the protein differ in their characteristics indicates that exit from the Golgi complex is linked to this change in the properties of caveolin. Exit from the Golgi complex is accelerated by the addition of cholesterol³⁰ and inhibited after glycosphingolipid depletion³¹. It is also associated with masking of specific caveolin epitopes (as shown using antibodies that recognize the Golgi pool, but not the plasma membrane pool, unless cholesterol is removed from the plasma membrane)³⁰. Light microscopy using CAV1-GFP (green fluorescent protein) revealed that defined quanta of caveolin form in the Golgi complex and are transported directly to the plasma membrane³². The quantal size is similar to that of surface caveolae, which indicates that exit from the Golgi is associated with caveolin assembly (oligomerization and association with cholesterol and glycosphingolipid-rich lipid-raft domains) to form a mature caveola-like exocytic structure (we propose the term 'exocytic caveolar carrier'), which is destined for the plasma membrane (FIG. 2). Whether this carrier is made up of only a mature caveola or has other non-caveolar membrane associated with it is as yet unknown.

How the assembly of a caveolin-enriched domain is linked to the budding of a mature caveolar carrier at this specific stage of the secretory pathway is not yet clear. However, the domain-induced budding model for lipid-raft-associated vesicle formation implies that glycosphingolipids and cholesterol might well be crucial factors in this process²³. Budding in this model is driven by the energy derived from the increase in the line tension that emanates from the increasing size of the lipid-raft domain that forms the caveolar carrier. A growing (phase-separated) domain will eventually reach a critical size, beyond which budding becomes energetically favourable³³. In line with this model, caveolar carriers would be formed late in the secretory pathway, presumably in the late Golgi compartments in which glycosphingolipids and cholesterol are present at high levels, rather than immediately after the synthesis of caveolin in the ER. However, whether induction of the budding of the carriers involves further regulation, such as phosphorylation, is still unclear³⁴.

Post-Golgi trafficking of caveolins. Some of the other components that are associated with the formation of caveolar carriers during Golgi exit (and/or caveolin post-Golgi

Box 1 | Defining caveolae

One problem in the field has been laxity concerning the definition of caveolae. We would like to restrict the term caveolae to invaginations of the plasma membrane with a diameter of 60–80 nm. These invaginations are formed by the polymerization of caveolins and contain a subset of lipid-raft components, including cholesterol and sphingolipids. Caveolae might exist as single pits or can form a cluster of caveolae with non-caveolar membrane between the pits (FIG. 2). Clathrin-coated pits can even be observed in membrane continuity with caveolae within these clusters⁵⁹. Caveolae can flatten out into the plasma membrane, thereby losing their caveolar identity. The term flat caveolae is a misnomer that only caused confusion in the field.

Signal recognition particle (SRP). A complex of polypeptides and RNA involved in synthesis of proteins on membrane-bound ribosomes of the ER. SRP interaction with a specific signal on the nascent polypeptide dictates co-translational insertion of the protein into the ER.

Detergent-resistant membrane (DRM). DRM fractions remain insoluble after cold Triton X-100 extraction. This is a crude biochemical measure for lipid-raft association.

Exocytic caveolar carrier
A carrier produced in the Golgi that resembles a fully formed caveola in caveolin density. We suggest the terms endocytic caveolar carrier and recycling caveolar carrier for budded caveolae or caveolae recycling back to the cell surface.

transport) are also starting to be elucidated. The SNARE protein **syntaxin-6** is involved in the delivery of CAV1, glycosylphosphatidylinositol (GPI)-anchored proteins and the ganglioside GM1 to the plasma membrane³⁵. However, CAV1 is not required for efficient plasma membrane delivery of GPI-anchored proteins³⁶. By contrast, **dysferlin**³⁷, the angiotensin receptor³⁸, the insulin receptor³⁹ and the stretch-activated channel short transient receptor potential channel-1 (**TRPC1**)^{40,41} seem to depend on caveolin for efficient surface delivery. These proteins have been shown to be inefficiently transported to the plasma membrane in cells, which either lack caveolins, or which express mutant Golgi-accumulated forms of caveolins. When examined in detail, at least some of these proteins did not localize to caveolae at the cell surface³⁸. Whether these proteins are transported to the plasma membrane in caveolar carriers or through other exocytic pathways but in a caveolin-dependent manner (FIG. 2) awaits further detailed characterization. In epithelial cells, CAV1 and CAV2 are targeted to the basolateral surface and form caveolae, whereas caveolae are not normally observed on the apical surface where only CAV1 is found^{42,43}. CAV1 presumably travels as a cargo protein from the Golgi to the apical surface in distinct, non-caveolar-carriers, which are not yet defined. A more general question is whether caveolae or caveolar carriers only incorporate cargo that is bound to the outer or inner leaflet, or whether they can also accommodate transmembrane proteins without perturbing the caveolin polymeric network in the lipid-raft cluster.

Box 2 | Caveolae in disease

Although the exact physiological roles of caveolae continue to be a matter of some debate, there is strong evidence for an association of caveolar dysfunction with human disease⁷. Caveolin-1 (CAV1)-null cells show increased proliferation, and loss of CAV1 accelerates tumourigenesis^{122,123}. In some breast cancers, CAV1 is downregulated, and a number of sporadic mutations in CAV1 have been detected in samples of human breast cancer^{124–126}, correlating specifically with oestrogen-receptor- α -positive status¹²⁷. In some tumours, CAV1 can also promote tumour survival and growth¹²⁸, and CAV1 has been associated with the progression of prostate carcinoma^{129,130}.

CAV3, the muscle-specific caveolin isoform, is also strongly linked to disease. Many mutations in CAV3 have now been described, and the mutant proteins are associated with a number of human muscle disorders including limb girdle muscular dystrophy and rippling muscle disease^{131–134}. The mutant CAV3 proteins often show mislocalization to the Golgi complex¹³⁵ and cause a reduction in the surface levels of CAV3 and other muscle plasma membrane proteins, such as the membrane-repair protein dysferlin^{37,136}.

Endocytosis of caveolae

Caveolae at the cell surface form a stable functional unit that is generated by oligomerized caveolin and associated proteins and lipids. Fluorescence recovery after photobleaching (FRAP) studies have shown that caveolins in caveolae are relatively immobile, but mobility can significantly increase upon cholesterol depletion^{44,45}. This finding has been confirmed by fusing cells with differently coloured caveolins; the caveolae remain red or green and do not form mixed structures³².

The caveolar unit is also maintained as a stable structure upon endocytosis⁴⁴. Budded caveolae, here known as endocytic caveolar carriers, can fuse with the caveosome in a **RAB5**-independent manner (BOX 3; FIG. 3), or with the early endosome in a RAB5-dependent manner. The caveolar unit can then be recycled back to the plasma membrane for reuse while maintaining the tight association of caveolin oligomers in a stable unit⁴⁴. Caveolar carriers can also fuse back to the plasma membrane without the involvement of an intermediate station, a process that is also subject to tight regulation²¹. In view of the apparent stability of the caveolar unit, an interesting question is how fusion with the plasma membrane or with caveosomes is achieved, and whether this requires partial disassembly of the caveolar coat. Furthermore, it is unclear whether caveolae always bud as single units, as shown in recent ultrastructural studies⁴⁶, or whether they can also detach from the membrane as part of larger domains with one or more caveolae connected to non-caveolar membrane.

Caveolae remain for long periods at the plasma membrane⁴⁶, but their internalization can be stimulated by various agents. These include the SV40 virus, which uses caveolae for entry into cells and stimulates caveolar budding^{8,47}, as well as sterols and glycosphingolipids⁴⁸. Despite the different agents used to stimulate caveola internalization, there are common mechanisms involved in these pathways with a crucial role for **dynamitin**^{49–51}, Src kinases, protein kinase C (PKC) and actin recruitment^{8,47,48}. Also other clathrin- and caveolin-independent, lipid-raft endocytosis pathways have been identified that differ from the caveolar pathway in their characteristics^{31,46,52,53}. This complicates the interpretation of studies of non-clathrin-mediated endocytic routes. Downregulation of CAV1 and the resulting loss of caveolae does not block the endocytosis of many lipid-raft markers — including cholera toxin, which binds to GM1 (REFS 53,54) — highlighting the multiple endocytic mechanisms that are involved in the internalization of these markers.

Caveolae and cell adhesion. A large-scale RNA interference screen examining the role of specific kinases in caveolar endocytosis has provided key insights into this process and its regulation⁵⁵. The results not only present a new global view of the interplay between different endocytic pathways but also provide fascinating insights into the possible function of caveolar endocytosis. For example, several kinases that regulate SV40 endocytosis are associated with the regulation of cell adhesions, focal contacts and extracellular matrix (ECM) interactions⁵⁵.




Figure 2 | Caveolin biosynthesis and trafficking to the plasma membrane. Caveolin is synthesized in the endoplasmic reticulum (ER) and is then transported to the Golgi complex as detergent-soluble oligomers. When caveolin exits the Golgi complex, the oligomers associate with glycosphingolipid- and cholesterol-enriched lipid-raft domains, as judged by detergent-resistant membrane (DRM) association. Caveolar carriers that contain quanta of caveolin are produced and fuse directly with the plasma membrane in a process that requires the SNARE protein syntaxin-6. In addition, we propose that caveolin can be transported to the plasma membrane by other carriers. Many caveolin mutants that have been associated with disease conditions accumulate in the Golgi complex and are not transported to the plasma membrane efficiently¹³⁵ (as indicated by mutant caveolin (red) in the figure). Expression of these mutants seems to cause a more complete block in Golgi exit, and be more harmful, than total loss of caveolin^{37,139,140}. One explanation for this is that proteins that exit the Golgi through the caveolar pathway are facilitated by caveolin, but can also follow other pathways, presumably less efficiently, if caveolin is absent. However, caveolin mutants (which cannot form caveolar carriers) perturb the exit pathway and trap their cargo in the Golgi complex so cargo are unable to 'escape' through other routes.

Other studies strengthen the link between caveolar endocytosis and cell adhesion. Glycosphingolipids, which stimulate caveolar endocytosis, cause clustering and internalization of integrins through a pathway that involves caveolae; this pathway is inhibited by cholesterol perturbation, CAV1 knockdown or dominant-negative mutants of dynamin and is dependent on Src kinase and PKC⁵⁶. **Fibronectin** internalization and degradation is also inhibited by knockdown of CAV1 (REF. 57).

Caveolar endocytosis also seems to be important under conditions in which cells lose adhesion to their substratum⁵⁸. The detachment of fibroblasts triggers internalization of the ganglioside GM1 from the plasma membrane in a caveolin-dependent manner. The internalization of GM1 causes loss of **Rac1** from the plasma membrane and thereby suppression of Rac1 activation. During the detachment process, Tyr14-phosphorylated CAV1 is localized in caveolae. This phosphorylation event is required for endocytosis — as shown using *Cav1*^{-/-} cells in which wild-type CAV1 or a non-phosphorylated CAV1 mutant had been reintroduced. The striking, rapid removal of GM1 and other lipid-raft components

from the cell surface due to internalization of caveolae, as proposed, is surprising in view of the limited enrichment of GM1 in caveolae under control conditions, as shown using both morphological and biochemical techniques^{22,59,60}. It has been proposed that CAV1 is required for the coordination of the detachment stimulus, which induces a massive lipid-raft clustering process (possibly macropinocytosis⁶¹), and leads to domain-induced inward budding of lipid-raft components and proteins, followed by internalization. Taken together these studies show strong links between caveolar endocytosis and the regulation of cell adhesion. But how these findings relate to the apparently normal development of *Cav1*-null mice, which completely lack caveolae, is still unclear.

Other roles for caveolar endocytosis. Endocytosis through caveolae has been extensively studied in endothelial cells but still remains a matter of much debate. Caveolae are particularly abundant in vascular endothelial cells *in vivo*. Endocytosis and transcytosis through caveolae have been claimed to be involved in the transcellular transport of albumin^{62,63}. However, the levels of tissue albumin seem normal in *Cav1*^{-/-} mice¹¹. In fact, loss of caveolae *in vivo* through small interfering (si)RNA-mediated downregulation of CAV1 resulted in an increase in vascular permeability to albumin⁶⁴. Ultrastructural studies indicate that the increased vascular permeability to albumin might be a result of the dilation of interendothelial junctions⁶⁴, which is consistent with another study showing higher transvascular transport of solutes in *Cav1*-null mice⁶⁵. This study shows similar passive transport across the endothelial monolayer in the *Cav1*-null and wild-type mice, arguing against a significant role for caveolar transcytosis in transport across the endothelium *in vivo*⁶⁵. A clear-cut role for caveolae in this process is not proven at present, and might be complicated by the effects of CAV1 loss on the activity of endothelial nitric-oxide synthase (**eNOS**) and the resulting changes in vascular permeability⁶⁶. In addition, as with other ligands and in other cell types, the dependence of albumin endocytosis on caveolae is not absolute and it cannot be excluded that other pathways take over in *Cav1*-null cells.

Other diverse roles for caveolar internalization have been described, for example, the endocytosis of epidermal growth factor receptor (EGFR) and transforming growth factor receptor (TGFR)^{57,68}. Again, these and other studies remain to be confirmed in view of the evidence for multiple clathrin-independent endocytosis pathways³¹.

Caveolae and large-scale membrane changes

Small viruses such as SV40 and polyoma virus (of approximately 45-nm diameter) enter cells through caveolae^{8,69,70}, and it has been suggested that a number of larger pathogens — which are far too large for endocytosis by a single caveola — also make use of caveolae as an endocytic vehicle. By exploiting this entry pathway, pathogens avoid degradation in lysosomes. Such pathogens include **FimH**-expressing *Escherichia coli*⁹ and *E. coli* K1 (REF. 71), *Pseudomonas aeruginosa*⁷² and *Porphyromonas gingivalis*⁷³. By exploiting this entry

SNAREs

(Soluble *N*-ethylmaleimide-sensitive factor attachment-protein receptors). A protein family that consists of a cognate group of integral and peripheral membrane proteins that are required for bilayer recognition and fusion during membrane trafficking.

Caveosome

A neutral pH endosomal compartment that lacks classic endosomal markers but contains markers that are internalized through caveolae.

Box 3 | Features and properties of the caveosome

The caveosome was identified by Helenius and Pelkmans as a distinct endosomal compartment that fuses with SV40-containing caveolae^{8,44,47} (FIG. 3). This compartment has specific properties:

- The caveosome is negative for markers of endosomes on the clathrin-mediated endocytic pathway, such as early endosomal protein-1 (EEA1) and internalized transferrin.
- The caveosome has a neutral luminal pH, which is in contrast to early endosomes, late endosomes and lysosomes, which have an acidic pH.
- The caveosome has some of the properties of a sorting compartment. For example, cholera toxin within the caveosome is sorted to carriers that are destined for the Golgi complex, whereas caveolin-1 (CAV1) returns to the plasma membrane. SV40 exits from the caveosome and moves along microtubule-dependent carriers to the endoplasmic reticulum.

It should be noted that not all CAV1-labelled vacuolar structures within the cell are caveosomes, as has been assumed in some electron-microscopy studies. Many cells have caveola-covered structures that seem to be intracellular vacuoles, but are actually connected to the plasma membrane out of the plane of section (see, for example, FIG. 1). An extreme example of this unusual plasma membrane organization is seen in adipocytes¹³⁷.

pathway, pathogens avoid degradation in lysosomes. The bacteria do not use caveolae as such, but they seem to be internalized in caveola-rich areas of the cell⁷⁴ through a non-clathrin cholesterol-dependent pathway, and often locate to caveolin-containing vacuoles within the cell. Although these bacteria are not internalized within caveolae, and the apical surface of epithelia where invasion often occurs lacks caveolae^{42,43}, several studies have shown an effect of caveolin downregulation on the entry process (see, for example, REF. 73).

How does loss of CAV1 affect the invasion process? The gross membrane rearrangements that are required to accommodate the invading pathogens might require coordination of membrane and cytoskeletal changes that are consistent with a general role for caveolin in these processes, rather than as an endocytic carrier. For example, group A streptococci bind to the ECM protein fibronectin, which in turn binds to integrins⁷⁴. These interactions then trigger the uptake of streptococci by large, encapsulating invaginations. This process resembles the dramatic internalization of ganglioside-rich domains that occurs after breaking focal adhesions⁵⁸. Both processes involve caveolins and could be shaped by massive lipid-raft clustering.

Another large-scale invagination process in which caveolae have been implicated is the transcellular migration of lymphocytes across the endothelium⁷⁵. CAV1 downregulation partially inhibits this process. Clustering of the endothelial adhesion molecule intercellular adhesion molecule-1 (ICAM1) was proposed to trigger transcytosis through caveolae; however, the role for caveolae in ICAM1 endocytosis remains controversial⁷⁶. Caveolins have also been implicated in phagocytosis⁷⁷ and cell motility⁷⁸. The latter could involve regulation of the focal-adhesion machinery and/or regulation of endocytic events. As will be discussed further below, caveolae might be well-equipped to sense changes in the membrane that are required for these large-scale surface rearrangements.

Caveolae and signalling

The number of papers linking caveolae to specific signalling pathways has grown exponentially. Are caveolae the universal signalling regulators originally proposed? If so, how do cells regulate signalling in the absence of caveolae, both in those cells that naturally lack caveolae (lymphocytes, for example) or in genetically modified systems (such as *Cav1*^{-/-} mice)? Are the scores of proteins that immunoprecipitate with CAV1 — including receptors and downstream signalling components, channels, structural proteins and enzymes — actually concentrated in caveolae? Purification of caveolae by immuno-isolation techniques, and analysis by immuno-electron microscopy have rarely supported these claims (see, for example, REF. 79) and in the absence of functional data these data might have to be re-examined (BOX 4). There are so many conflicting claims that scepticism should be maintained until further supporting data become available.

Universal signalling regulators? A key feature of the signalling model is the interaction between the conserved scaffolding domain of caveolin (amino acids 82–101 of CAV1) and signalling proteins⁸⁰. However, this domain also has membrane-binding activity^{81,82}, and might even insert into the caveolar membrane⁸³, possibly as an in-plane amphipathic helix²⁷. Antibodies against this region do not label caveolin in caveolae at the cell surface, but do label Golgi-based caveolin³⁰, possibly reflecting masking of the scaffolding domain during caveola formation. The caveolin-signalling hypothesis would predict that, in cells lacking caveolae, the distribution of signalling proteins would be affected. However, this is not the case; siRNA-mediated knockdown of CAV1 in aortic endothelial cells did not affect lipid-raft association or targeting of any of the signalling proteins that were tested in recent studies⁸⁴.

How can interactions between the scaffolding domain of caveolin and signalling proteins be explained? These interactions have been well-characterized in the case of eNOS and particularly through the use of peptides from the CAV1-scaffolding domain⁸⁵. One possibility is that the non-caveolar pool of CAV1, at the cell surface⁸⁶ or in the secretory pathway³⁸, is responsible for some of these interactions. In support of this, transgenic overexpression of CAV1 inhibits eNOS activation without increasing caveola formation⁸⁷. This hypothesis could explain other reported interactions between expressed caveolins and signalling molecules, in situations where their presence in caveolae is not observed. Caveolae could provide a reservoir of caveolin molecules that are only released under specific cellular conditions. This hypothesis must be examined carefully in future studies.

These questions aside, there is evidence of a role for caveolins in specific signalling pathways that is strongly supported by functional studies. In addition to a role in eNOS regulation^{88,89}, several other pathways show a crucial dependence on caveolins^{84,90,91}. Caveolae have also been implicated in the regulation of channels and in calcium signalling^{92,93}, although the precise role of caveolae or non-caveolar caveolins is not yet clear.




Figure 3 | Caveola endocytosis. Caveolae at the cell surface can bud into the cell carrying cholera-toxin-binding subunit (CTxB) and SV40 in a process that is regulated by dynamin, protein kinase C and tyrosine kinases, such as Src kinases. Caveolae bud off to form endocytic caveolar carriers that fuse with the caveosome or with the early endosome, or can fuse back to the plasma membrane without the involvement of an endosomal intermediate^{8,44,141}. From the caveosome, SV40 is transported to the endoplasmic reticulum (ER) and recycling endocytic caveolar carriers carry caveolin back to the plasma membrane. CTxB is transported to the Golgi complex, possibly through early endosomes. Note that in cells with or without caveolae there is at least one other main endocytic pathway that involves clathrin- (and caveolin)-independent carriers (CLICs), which can endocytose the same surface markers into endosomes⁴⁶. Glycosylphosphatidylinositol (GPI)-anchored proteins are predominantly internalized through this pathway.

Caveolins and lipid regulation

Caveolae are extremely abundant in adipocytes, and increasing evidence has linked caveolae to lipid regulation in this specialized lipid-storage cell as well as in other cell types. CAV1 interacts with cholesterol, binds to fatty acids⁹⁴ and associates with lipid droplets (the site of cellular lipid storage^{95,96}) under specific cellular conditions in cultured cells and *in vivo*^{97–99}.

To examine the role of caveolins in lipid regulation, researchers have studied the effect of *Cav1* expression in cells with low endogenous caveolin levels, as well as looking at the effect of caveolin mutants. Expression of *Cav1* facilitates the uptake of fatty acids into cells¹⁰⁰, and increases the levels of free cholesterol and cholesterol export^{101,102}. Studies of fibroblasts and peritoneal macrophages from *Cav1*-null mice showed a slight decrease in cholesterol synthesis and increased esterification in both cell types¹⁰³. A caveolin truncation mutant constitutively associated with lipid droplets and perturbed cellular lipid regulation, which decreased surface levels of free cholesterol and increased neutral lipid storage in lipid droplets^{99,104}. Taken together, these studies all point to a role for CAV1 in sterol and fatty-acid regulation, both at the plasma membrane and possibly in lipid droplets; however, the detailed mechanisms are still unclear. The importance of this process *in vivo* is emphasized by the requirement for CAV1 for efficient liver regeneration and mouse survival after partial hepatectomy¹⁰⁵. The lack of CAV1 caused a dramatic reduction in lipid-droplet formation

during the liver-regeneration process. Survival could be dramatically increased by providing glucose as an alternative energy source¹⁰⁵.

Lipid sensing and storage. Adipocytes have attracted considerable interest in view of the density of caveolae in these cells and their vital role in lipid regulation. *Cav1*^{−/−} mice show decreased adiposity and are resistant to diet-induced obesity¹⁰⁶. Caveolae at the cell surface of adipocytes have been implicated in triacylglycerol synthesis, but there is also evidence for the redistribution of CAV1 from the cell surface to lipid droplets upon treatment with cholesterol¹⁰⁷ and upon stimulation of lipolysis⁹⁸. The cholesterol-stimulated association of CAV1 with lipid droplets is inhibited by agents that perturb caveola internalization whereas caveola budding is stimulated by cholesterol¹⁰⁷. These data indicate a novel, as-yet-uncharacterized, pathway by which CAV1 reaches the lipid droplet, and also link the endocytic role of caveolae with lipid sensing and storage. The intermediate stations on this pathway are unknown, although it is clear that caveolae cannot directly fuse with the lipid droplet, which is enclosed by a phospholipid monolayer. Rather, caveolin might flip from the cytoplasmic leaflet of one membrane into the monolayer. The functional importance of this pathway is unclear; however, the finding that adipocytes from *Cav1*^{−/−} mice show decreased levels of free cholesterol in the lipid droplet¹⁰⁷ indicates an important role in regulating cholesterol trafficking to, or association with, the lipid droplet.

Lipid droplet

A lipid-storage organelle that comprises a core of triacylglycerol and/or cholesterol esters surrounded by a phospholipid monolayer.

Box 4 | Approaches to functionally characterize caveolae

Several functions have been assigned to caveolae and caveolins but many of these functions must be critically reassessed. In many cases, numerous strategies were used for studying these structures and all of them contributed to the conclusion of caveolar involvement in a specific process and/or the association of proteins with caveolae. These approaches include the following:

- cholesterol depletion
- fractionation using detergents or detergent-free methods
- immunofluorescence
- immunoprecipitation.

There are problems associated with each approach and even their combined use can provide misleading results. First, cholesterol depletion is not specific to caveolae; even in cells that lack caveolae, cholesterol depletion affects many cellular processes. Second, detergent-insoluble preparation and detergent-free methods of purification do not yield caveolar preparations¹³⁸. Studies of caveolae have also been confounded by the small size of caveolae and the resolution of light microscopy. Numerous studies have provided evidence for the colocalization of particular proteins with caveolin proteins using light microscopy, but inspection of the images shows only partial overlap, consistent simply with the observation that two surface proteins are present on the plasma membrane.

The immunoprecipitation of caveolins with hundreds of different proteins is also puzzling. In some cases, solubilization conditions are unlikely to have caused the dissociation of detergent-resistant membranes (for example, immunoprecipitation from detergent-resistant membranes without further solubilization or with mild detergents). The vast array of proteins that have been shown to immunoprecipitate with caveolin must be considered with a measure of scepticism if the results are not confirmed by independent techniques. In several cases, the association of proteins with caveolae have not been confirmed by immuno-electron microscopy or immuno-isolation; however, an interesting possibility is that non-caveolar caveolin might explain some of these results. As outlined in the main text, the study of caveolin-null animals and the small interfering (si)RNA-based knockdown of caveolins has started to provide exciting new insights into the involvement of caveolae and caveolins in specific cellular processes.

Caveolae as lipid-raft stores. One astounding property of caveolae is the stability of their lipid-protein assembly. Caveolae seem to be largely static surface structures at the plasma membrane, and so far we only know of viruses, exogenous gangliosides or cholesterol that can trigger their internalization. In some cells, such as endothelial cells, caveolae represent more than one-third of the plasma membrane surface area. Some of these caveolae fold to form 'bunches of grapes' of connected caveolae (FIG. 1), which prompted Tomas Landh to postulate that these structures could represent cubic membrane infoldings that can fold and unfold with little energy input¹⁰⁸.

Why do some cell types have so many caveolae attached to their plasma membranes? Could caveolae function as stores for sphingolipids and cholesterol? These stores could ensure a supply of lipid-raft components to the cell when needed. Caveolae could also be used as a device to regulate plasma membrane surface area. Some cell types have such a high density of caveolae that flattening of the caveolae could contribute significantly to an increase in the accessible surface area of the cell.

We therefore postulate that mammalian cells could use two different organelles to store lipids: lipid droplets and caveolae. If this is the case, regulatory circuits that coordinate their functions in different cell types must be in place.

Caveolae as mechanosensors

One of the most exciting insights into caveola function that has been derived from the use of caveolin-null mice and knockdown systems is the demonstration of a role for caveolae in mechanosensation (for a review see REF. 109). The ability to sense membrane forces might underlie many of the functions of caveolae in diverse cell types, including endothelial and smooth-muscle cells.

Flow sensors in endothelial cells. The surface of endothelial cells is sensitive to changes in hydrostatic pressure and shear stresses. Chronic exposure to shear stress resulted in increased plasma membrane levels of CAV1 due to redistribution of CAV1 from the Golgi complex to the plasma membrane, which caused an increased surface density of caveolae^{110,111}. These changes are accompanied by increased mechanosensitivity and activation of specific signalling pathways (including eNOS and mitogen-activated protein kinase (MAPK)) and increased phosphorylation of CAV1. Subsequent studies confirmed and extended these findings and linked integrins to CAV1 phosphorylation¹¹².

The role of caveolae in mechanosensing is strengthened by recent *in vivo* studies using *Cav1*^{-/-} mice¹¹³. *Cav1*-knockout mice were crossed with mice expressing CAV1 under an endothelium-specific promoter to assess the precise role of endothelial caveolae *in vivo*. These studies showed a role for CAV1 and caveolae in mechanotransduction and remodelling of blood vessels. *Cav1*-null mice showed defects in chronic flow-dependent remodelling, and in acute flow-dependent dilation; both effects were rescued by CAV1 re-expression in the endothelium, which indicates that caveolin/caveolae might represent a flow sensor.

The endothelial phenotype that has been observed after chronic shear stress is similar to that observed in eNOS-deficient mice in response to flow¹¹⁴. In fact, a decrease in eNOS activation was observed in the *Cav1*-null mice¹¹³. This observation elegantly demonstrates that coupling of the flow stimulus to eNOS activation is lost in the absence of CAV1 and caveolae. This model might seem paradoxical in view of the postulated role of caveolae in the negative regulation of eNOS, and the increased nitric oxide production that has been reported in *Cav1*^{-/-} mice. Based on the above findings, Yu and colleagues proposed that the coupling of flow to eNOS activation might require caveolae, whereas activation of eNOS by agonists is a different process that might be hyperstimulated in the absence of caveolae owing to the loss of the inhibitory influence of CAV1 on eNOS function¹¹³.

Stretch-induced cell-cycle progression. Ongoing studies of caveolae in smooth-muscle cells also support a role for caveolae in mechanosensing. Smooth-muscle cells that have been subjected to cyclic stretch show rapid redistribution of CAV1 to focal contacts¹¹⁵. In control cells, stretch triggers cell-cycle progression through the phosphatidylinositol 3-kinase (PI3K)-AKT/protein kinase B (PKB) pathway, the MAPK ERK, c-Src, and integrins. This response is inhibited upon transient downregulation of CAV1 or in *Cav1*^{-/-} smooth-muscle

Mechanosensation

The sensing of mechanical stimuli, for example stretch or flow, by cells.

cells, which indicates an essential role for CAV1 in this process. This requirement was recapitulated in an *in vivo* system¹¹⁵.

Molecular mechanisms of mechanosensing. Caveolae have been shown to harbour G_q proteins that could be activated by unknown mechanotransducers¹¹⁶, as well as a sphingomyelinase that could produce ceramide from caveolar sphingomyelin stores in response to specific stimuli¹¹⁷. Integrin-mediated connectivity between the ECM and the cytoskeleton has also been implicated in mechanotransduction¹¹⁸. Caveolin and caveolae have been tightly connected to integrins and the focal-adhesion machinery. Caveolae can also be removed from the surface by endocytosis in response to specific stimuli, which provides a mechanism to regulate surface mechanosensing activity.

However, one question still remains. How might caveolar functions be regulated by mechanical strain? Could the specialized lipid composition of caveolae be involved? One possibility is that mechanical stress could cause changes in membrane elasticity¹¹⁹. This is determined by the lipid composition of the bilayer. Membrane elasticity regulates the hydrophobic coupling between a membrane-spanning protein and the surrounding bilayer. Changes in cholesterol concentration have been shown to have a specific role in modifying the elastic properties of this coupling¹²⁰. Therefore, it is not unreasonable to speculate that mechanosensitive cholesterol–protein couples could be part of the transducing machinery. Mechanical stress and the addition of lipids to cells cause a similar caveolar response. Only further work will clarify whether such mechanisms really are operating. Sens and Turner¹²¹ have recently analysed the issue of budded microdomains as tension regulators from the biophysical viewpoint. They came to the conclusion that caveolae could perform such a role by flattening out into the plasma membrane as a response to increases in surface tension.

Conclusions and perspectives

Caveolae are emerging as important surface-associated organelles that have vital functions in diverse cellular processes. Caveolae are formed by the polymerization of caveolins, which leads to the clustering and invagination of a subset of sphingolipid–cholesterol lipid rafts. In some cell types, considerable reservoirs of lipid-raft components are stored in these surface invaginations. These structures are stable; the bulk of the caveolar caveolins can only be released by dissociation of the caveola structure. But how this disassembly process is regulated is not known.

Nevertheless, there is evidence for a second pool of non-caveolar caveolins in focal adhesions. Detachment of cells from the substratum by the breaking of their focal adhesions leads to massive caveolin-dependent endocytosis. Bacteria might use similar mechanisms, which are possibly also triggered by integrin activation. Other stimuli for caveolar endocytosis include viruses, exogenous glycosphingolipids or addition of cholesterol. How internalization is regulated physiologically and for what purpose is not known; however, caveolar endocytosis seems to be linked to the regulation of the cellular lipid balance.

The most intriguing property of the caveolin/caveolae system is its involvement in mechanosensing. Many of the apparently unrelated functions of caveolae might be related to the ability of caveolae to sense changes in membrane tension or to sense other changes in the physical properties of the plasma membrane (for example, those caused by lipid loading). In this model, caveolae would coordinate physical membrane changes with intracellular signal transduction pathways. How caveolae are structured to carry out these different functions remains to be explored. However, the tools are now there to resolve how caveolae function at the molecular level in cell physiology and pathology.

- Palade, G.E. Fine structure of blood capillaries. *J. Appl. Phys.* **24**, 1424 (1953).
- Yamada, E. The fine structures of the gall bladder epithelium of the mouse. *J. Biophys. Biochem. Cytol.* **1**, 445–458 (1955).
- Stan, R.V. Structure of caveolae. *Biochim. Biophys. Acta* **1746**, 334–348 (2005).
- Stan, R.V., Tkachenko, E. & Niesman, I.R. PV1 is a key structural component for the formation of the stomatal and fenestral diaphragms. *Mol. Biol. Cell* **15**, 3615–3630 (2004).
- Lisanti, M.P. *et al.* Caveolae, transmembrane signalling and cellular transformation. *Mol. Membr. Biol.* **12**, 121–124 (1995).
- Kurzchalia, T.V. & Parton, R.G. Membrane microdomains and caveolae. *Curr. Opin. Cell Biol.* **11**, 424–431 (1999).
- Williams, T.M. & Lisanti, M.P. Caveolin-1 in oncogenic transformation, cancer, and metastasis. *Am. J. Physiol. Cell Physiol.* **288**, C494–C506 (2005).
- Pelkmans, L., Kartenbeck, J. & Helenius, A. Caveolar endocytosis of simian virus 40 reveals a new two-step vesicular-transport pathway to the ER. *Nature Cell Biol.* **3**, 473–483 (2001).
- Shin, J.S., Gao, Z. & Abraham, S.N. Involvement of cellular caveolae in bacterial entry into mast cells. *Science* **289**, 785–788 (2000).
- Fra, A.M., Williamson, E., Simons, K. & Parton, R.G. *De novo* formation of caveolae in lymphocytes by expression of VIP21-caveolin. *Proc. Natl Acad. Sci. USA* **92**, 8655–8659 (1995).
- Drab, M. *et al.* Loss of caveolae, vascular dysfunction, and pulmonary defects in Caveolin-1 gene-disrupted mice. *Science* **293**, 2449–2452 (2001).
- Galbati, F. *et al.* Caveolin-3 null mice show a loss of caveolae, changes in the microdomain distribution of the dystrophin–glycoprotein complex, and t-tubule abnormalities. *J. Biol. Chem.* **276**, 21425–21433 (2001).
- Tang, Z. *et al.* Molecular cloning of caveolin-3, a novel member of the caveolin gene family expressed predominantly in muscle. *J. Biol. Chem.* **271**, 2255–2261 (1996).
- Way, M. & Parton, R.G. M-caveolin, a muscle-specific caveolin-related protein. *FEBS Lett.* **376**, 108–112 (1995).
- Razani, B. *et al.* Caveolin-2-deficient mice show evidence of severe pulmonary dysfunction without disruption of caveolae. *Mol. Cell. Biol.* **22**, 2329–2344 (2002).
- Sowa, G., Pypaert, M., Fulton, D. & Sessa, W.C. The phosphorylation of caveolin-2 on serines 23 and 36 modulates caveolin-1-dependent caveolae formation. *Proc. Natl Acad. Sci. USA* **100**, 6511–6516 (2003).
- Lahtinen, U., Honsho, M., Parton, R.G., Simons, K. & Verkade, P. Involvement of caveolin-2 in caveolar biogenesis in MDCK cells. *FEBS Lett.* **538**, 85–88 (2003).
- Murata, M. *et al.* VIP21/caveolin is a cholesterol-binding protein. *Proc. Natl Acad. Sci. USA* **92**, 10339–10343 (1995).
- Dietzen, D.J., Hastings, W.R. & Lublin, D.M. Caveolin is palmitoylated on multiple cysteine residues. Palmitoylation is not necessary for localization of caveolin to caveolae. *J. Biol. Chem.* **270**, 6838–6842 (1995).
- Rothberg, K.G. *et al.* Caveolin, a protein component of caveolae membrane coats. *Cell* **68**, 673–682 (1992).
- Pelkmans, L. & Zerial, M. Kinase-regulated quantal assemblies and kiss-and-run recycling of caveolae. *Nature* **436**, 128–133 (2005).
- Ortengren, U. *et al.* Lipids and glycosphingolipids in caveolae and surrounding plasma membrane of primary rat adipocytes. *Eur. J. Biochem.* **271**, 2028–2036 (2004).
- Quantification of lipid levels within caveolae shows that glycosphingolipids and cholesterol are concentrated more within caveolae compared with the surrounding plasma membrane. Higher packing of total lipids is observed in caveolae compared with the surrounding plasma membrane.**
- Schuck, S. & Simons, K. Polarized sorting in epithelial cells: raft clustering and the biogenesis of the apical membrane. *J. Cell Sci.* **117**, 5955–5964 (2004).
- Simons, K. & Vaz, W.L. Model systems, lipid rafts, and cell membranes. *Annu. Rev. Biophys. Biomol. Struct.* **33**, 269–295 (2004).
- An extensive review of the data supporting lipid-based microdomain formation in model systems and in cell membranes.**
- Hancock, J.F. Lipid rafts: contentious only from simplistic standpoints. *Nature Rev. Mol. Cell Biol.* **7**, 456–462 (2006).

26. Lipardi, C. *et al.* Caveolin transfection results in caveolae formation but not apical sorting of glycosylphosphatidylinositol (GPI)-anchored proteins in epithelial cells. *J. Cell Biol.* **140**, 617–626 (1998).
27. Parton, R.G., Hanzal-Bayer, M. & Hancock, J.F. Biogenesis of caveolae: a structural model for caveolin-induced domain formation. *J. Cell Sci.* **119**, 787–796 (2006).
- Summary of current data on caveola biogenesis and a proposed model for the formation of caveolae through caveolin–lipid interactions.**
28. Monier, S. *et al.* VIP21–caveolin, a membrane protein constituent of the caveolar coat, oligomerizes *in vivo* and *in vitro*. *Mol. Biol. Cell* **6**, 911–927 (1995).
29. Monier, S., Dietzen, D.J., Hastings, W.R., Lublin, D.M. & Kurzchalia, T.V. Oligomerization of VIP21–caveolin *in vitro* is stabilized by long chain fatty acylation or cholesterol. *FEBS Lett.* **388**, 143–149 (1996).
30. Pol, A. *et al.* Cholesterol and fatty acids regulate dynamic caveolin trafficking through the Golgi complex and between the cell surface and lipid bodies. *Mol. Biol. Cell* **16**, 2091–2105 (2005).
31. Cheng, Z.J. *et al.* Distinct mechanisms of clathrin-independent endocytosis have unique sphingolipid requirements. *Mol. Biol. Cell* **17**, 3197–3210 (2006).
32. Tagawa, A. *et al.* Assembly and trafficking of caveolar domains in the cell: caveolae as stable, cargo-triggered, vesicular transporters. *J. Cell Biol.* **170**, 769–779 (2005).
- Real-time microscopy and cell-fusion experiments show that caveolae are remarkably stable with evidence for formation of 'caveolae' in the Golgi complex that then fuse directly with the plasma membrane.**
33. Lipowsky, R. Domain-induced budding of fluid membranes. *Biophys. J.* **64**, 1133–1138 (1993).
34. Bauer, M. & Pelkmans, L. A new paradigm for membrane-organizing and -shaping scaffolds. *FEBS Lett.* **580**, 5559–5564 (2006).
35. Choudhury, A., Marks, D.L., Proctor, K.M., Gould, G.W. & Pagano, R.E. Regulation of caveolar endocytosis by syntaxin 6-dependent delivery of membrane components to the cell surface. *Nature Cell Biol.* **8**, 317–328 (2006).
- Identifies syntaxin-6 as a regulator of CAV1, GPI-anchored proteins, and GM1 transport from Golgi to the plasma membrane and provides insights into the coupling of exocytosis and endocytosis of lipid-raft components.**
36. Manninen, A. *et al.* Caveolin-1 is not essential for biosynthetic apical membrane transport. *Mol. Cell Biol.* **25**, 10087–10096 (2005).
37. Hernandez-Deviez, D.J. *et al.* Aberrant dysferlin trafficking in cells lacking caveolin or expressing dystrophy mutants of caveolin-3. *Hum. Mol. Genet.* **15**, 129–142 (2006).
38. Wyse, B.D. *et al.* Caveolin interacts with the angiotensin II type 1 receptor during exocytic transport but not at the plasma membrane. *J. Biol. Chem.* **278**, 23738–23746 (2003).
39. Cohen, A.W. *et al.* Caveolin-1-deficient mice show insulin resistance and defective insulin receptor protein expression in adipose tissue. *Am. J. Physiol. Cell Physiol.* **285**, C222–C235 (2003).
40. Brazer, S.C., Singh, B.B., Liu, X., Swaim, W. & Ambudkar, I.S. Caveolin-1 contributes to assembly of store-operated Ca^{2+} influx channels by regulating plasma membrane localization of TRPC1. *J. Biol. Chem.* **278**, 27208–27215 (2003).
41. Maroto, R. *et al.* TRPC1 forms the stretch-activated cation channel in vertebrate cells. *Nature Cell Biol.* **7**, 179–185 (2005).
42. Scheiffele, P. *et al.* Caveolin-1 and -2 in the exocytic pathway of MDCK cells. *J. Cell Biol.* **140**, 795–806 (1998).
43. Verkade, P., Harder, T., Lafont, F. & Simons, K. Induction of caveolae in the apical plasma membrane of Madin–Darby canine kidney cells. *J. Cell Biol.* **148**, 727–739 (2000).
44. Pelkmans, L., Burli, T., Zerial, M. & Helenius, A. Caveolin-stabilized membrane domains as multifunctional transport and sorting devices in endocytic membrane traffic. *Cell* **118**, 767–780 (2004).
45. Thomsen, P., Roepstorff, K., Stahlhut, M. & van Deurs, B. Caveolae are highly immobile plasma membrane microdomains, which are not involved in constitutive endocytic trafficking. *Mol. Biol. Cell* **13**, 238–250 (2002).
46. Kirkham, M. *et al.* Ultrastructural identification of uncoated caveolin-independent early endocytic vesicles. *J. Cell Biol.* **168**, 465–476 (2005).
47. Pelkmans, L., Puntener, D. & Helenius, A. Local actin polymerization and dynamin recruitment in SV40-induced internalization of caveolae. *Science* **296**, 535–539 (2002).
48. Sharma, D.K. *et al.* Selective stimulation of caveolar endocytosis by glycosphingolipids and cholesterol. *Mol. Biol. Cell* **15**, 3114–3122 (2004).
49. Oh, P., McIntosh, D.P. & Schnitzer, J.E. Dynamin at the neck of caveolae mediates their budding to form transport vesicles by GTP-driven fission from the plasma membrane of endothelium. *J. Cell Biol.* **141**, 101–114 (1998).
50. Henley, J.R., Krueger, E.W., Oswald, B.J. & McNiven, M.A. Dynamin-mediated internalization of caveolae. *J. Cell Biol.* **141**, 85–99 (1998).
51. Yao, Q. *et al.* Caveolin-1 interacts directly with dynamin-2. *J. Mol. Biol.* **348**, 491–501 (2005).
52. Damm, E.M. *et al.* Clathrin- and caveolin-1-independent endocytosis: entry of simian virus 40 into cells devoid of caveolae. *J. Cell Biol.* **168**, 477–488 (2005).
53. Kirkham, M. & Parton, R.G. Clathrin-independent endocytosis: new insights into caveolae and non-caveolar lipid raft carriers. *Biochim. Biophys. Acta* **1745**, 273–286 (2005).
54. Nabi, I.R. & Le, P.U. Caveolae/raft-dependent endocytosis. *J. Cell Biol.* **161**, 675–677 (2003).
55. Pelkmans, L. *et al.* Genome-wide analysis of human kinases in clathrin- and caveolae/raft-mediated endocytosis. *Nature* **436**, 78–86 (2005).
- A high-throughput screen of kinases involved in endocytosis provides fascinating insights into the complex interplay between endocytosis and other cellular processes, such as cell adhesion and cell division.**
56. Sharma, D.K. *et al.* The glycosphingolipid, lactosylceramide, regulates β 1-integrin clustering and endocytosis. *Cancer Res.* **65**, 8233–8241 (2005).
- Evidence for glycosphingolipid-stimulated β 1-integrin internalization through a caveolae-mediated pathway.**
57. Sottile, J. & Chandler, J. Fibronectin matrix turnover occurs through a caveolin-1-dependent process. *Mol. Cell Biol.* **16**, 757–768 (2005).
58. del Pozo, M.A. *et al.* Phospho-caveolin-1 mediates integrin-regulated membrane domain internalization. *Nature Cell Biol.* **7**, 901–908 (2005).
- Demonstrates striking internalization of lipid-raft components upon cell detachment from the substratum through a phospho-CAV1-dependent pathway.**
59. Parton, R.G. Ultrastructural localization of gangliosides; GM1 is concentrated in caveolae. *J. Histochem. Cytochem.* **42**, 155–166 (1994).
60. Nichols, B.J. GM1-containing lipid rafts are depleted within clathrin-coated pits. *Curr. Biol.* **13**, 686–690 (2003).
61. Watarai, M., Makino, S., Fujii, Y., Okamoto, K. & Shirahata, T. Modulation of Brucella-induced macropinocytosis by lipid rafts mediates intracellular replication. *Cell Microbiol.* **4**, 341–355 (2002).
62. Minshall, R.D. *et al.* Endothelial cell-surface gp60 activates vesicle formation and trafficking via G(i)-coupled Src kinase signaling pathway. *J. Cell Biol.* **150**, 1057–1070 (2000).
63. Schubert, W. *et al.* Caveolin-deficient endothelial cells show defects in the uptake and transport of albumin *in vivo*. *J. Biol. Chem.* **276**, 48619–48622 (2001).
64. Miyawaki-Shimizu, K. *et al.* siRNA-induced caveolin-1 knockdown in mice increases lung vascular permeability via the junctional pathway. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **290**, L405–L413 (2006).
- siRNA-mediated downregulation of CAV1 *in vivo* causes a loss of endothelial caveolae and an increase in vascular hyperpermeability to albumin. Ultrastructural studies showed dilation of interendothelial junctions.**
65. Rosengren, B.I. *et al.* Transvascular protein transport in mice lacking endothelial caveolae. *Am. J. Physiol. Heart Circ. Physiol.* **291**, H1371–H1377 (2006).
- Evidence for a passive porous pathway for transport of albumin across the endothelium *in vivo* in both Cav1-knockout and wild-type mice, arguing against caveolae as a quantitatively important transport route.**
66. Schubert, W. *et al.* Microvascular hyperpermeability in caveolin-1 (–/–) knock-out mice. Treatment with a specific nitric-oxide synthase inhibitor, L-NAME, restores normal microvascular permeability in Cav1 null mice. *J. Biol. Chem.* **277**, 40091–40098 (2002).
67. Sigismund, S. *et al.* Clathrin-independent endocytosis of ubiquitinated cargos. *Proc. Natl Acad. Sci. USA* **102**, 2760–2765 (2005).
68. Di Guglielmo, G.M., Le Roy, C., Goodfellow, A.F. & Wrana, J.L. Distinct endocytic pathways regulate TGF- β receptor signalling and turnover. *Nature Cell Biol.* **5**, 410–421 (2003).
69. Anderson, H.A., Chen, Y. & Norkin, L.C. Bound simian virus 40 translocates to caveolin-enriched membrane domains, and its entry is inhibited by drugs that selectively disrupt caveolae. *Mol. Biol. Cell* **7**, 1825–1834 (1996).
70. Stang, E., Kartenbeck, J. & Parton, R.G. Major histocompatibility complex class I molecules mediate association of SV40 with caveolae. *Mol. Biol. Cell* **8**, 47–57 (1997).
71. Sukumaran, S.K., Quon, M.J. & Prasadara, N.V. *Escherichia coli* K1 internalization via caveolae requires caveolin-1 and protein kinase C α interaction in human brain microvascular endothelial cells. *J. Biol. Chem.* **277**, 50716–50724 (2002).
72. Zaas, D.W., Duncan, M.J., Li, G., Wright, J.R. & Abraham, S.N. *Pseudomonas* invasion of type I pneumocytes is dependent on the expression and phosphorylation of caveolin-2. *J. Biol. Chem.* **280**, 4864–4872 (2005).
73. Tamai, R., Asai, Y. & Ogawa, T. Requirement for intercellular adhesion molecule 1 and caveolae in invasion of human oral epithelial cells by *Porphyromonas gingivalis*. *Infect. Immun.* **73**, 6290–6298 (2005).
74. Rohde, M., Muller, E., Chhatwal, G.S. & Talay, S.R. Host cell caveolae act as an entry-port for group A streptococci. *Cell. Microbiol.* **5**, 323–342 (2003).
75. Millan, J. *et al.* Lymphocyte transcellular migration occurs through recruitment of endothelial ICAM-1 to caveola- and F-actin-rich domains. *Nature Cell Biol.* **8**, 113–123 (2006).
76. Muro, S. *et al.* A novel endocytic pathway induced by clustering endothelial ICAM-1 or PECAM-1. *J. Cell Sci.* **116**, 1599–1609 (2003).
77. Li, J. *et al.* Impaired phagocytosis in caveolin-1 deficient macrophages. *Cell Cycle* **4**, 1599–1607 (2005).
78. Beardsley, A. *et al.* Loss of caveolin-1 polarity impedes endothelial cell polarization and directional movement. *J. Biol. Chem.* **280**, 3541–3547 (2005).
- Provides evidence for a functional role for CAV1 in endothelial-cell motility. In actively migrating cells, caveolae are localized at the rear of the cell and excluded from the leading edge.**
79. Souto, R.P. *et al.* Immunopurification and characterization of rat adipocyte caveolae suggest their dissociation from insulin signaling. *J. Biol. Chem.* **278**, 18321–18329 (2003).
80. Li, S., Couet, J. & Lisanti, M.P. Src tyrosine kinases, G α subunits, and H-Ras share a common membrane-anchored scaffolding protein, caveolin. Caveolin binding negatively regulates the auto-activation of Src tyrosine kinases. *J. Biol. Chem.* **271**, 29182–29190 (1996).
81. Schlegel, A., Schwab, R.B., Scherer, P.E. & Lisanti, M.P. A role for the caveolin scaffolding domain in mediating the membrane attachment of caveolin-1. The caveolin scaffolding domain is both necessary and sufficient for membrane binding *in vitro*. *J. Biol. Chem.* **274**, 22660–22667 (1999).
82. Epan, R.M., Sayer, B.G. & Epan, R.F. Caveolin scaffolding region and cholesterol-rich domains in membranes. *J. Mol. Biol.* **345**, 339–350 (2005).
83. Arbuza, A. *et al.* Membrane binding of peptides containing both basic and aromatic residues. Experimental studies with peptides corresponding to the scaffolding region of caveolin and the effector region of MARCKS. *Biochemistry* **39**, 10330–10339 (2000).
84. Gonzalez, E., Nagiel, A., Lin, A.J., Golan, D.E. & Michel, T. Small interfering RNA-mediated down-regulation of caveolin-1 differentially modulates signaling pathways in endothelial cells. *J. Biol. Chem.* **279**, 40659–40669 (2004).
85. Bernatchez, P.N. *et al.* Dissecting the molecular control of endothelial NO synthase by caveolin-1 using cell-permeable peptides. *Proc. Natl Acad. Sci. USA* **102**, 761–766 (2005).
86. Wary, K.K., Mariotti, A., Zurzolo, C. & Giancotti, F.G. A requirement for caveolin-1 and associated kinase Fyn in integrin signaling and anchorage-dependent cell growth. *Cell* **94**, 625–634 (1998).

87. Bauer, P.M. *et al.* Endothelial-specific expression of caveolin-1 impairs microvascular permeability and angiogenesis. *Proc. Natl Acad. Sci. USA* **102**, 204–209 (2005).
Transgenic overexpression of CAV1 impairs eNOS activation and decreases VEGF-stimulated vascular permeability. CAV1 overexpression reduced VEGF-mediated angiogenesis after experimentally induced tissue ischaemia.
88. Bucci, M. *et al.* *In vivo* delivery of the caveolin-1 scaffolding domain inhibits nitric oxide synthesis and reduces inflammation. *Nature Med.* **6**, 1362–1367 (2000).
89. Garcia-Cardena, G. *et al.* Dissecting the interaction between nitric oxide synthase (NOS) and caveolin. Functional significance of the NOS caveolin binding domain *in vivo*. *J. Biol. Chem.* **272**, 25437–25440 (1997).
90. Reddy, M.A. *et al.* Key role of Src kinase in S100B-induced activation of the receptor for advanced glycation end products in vascular smooth muscle cells. *J. Biol. Chem.* **281**, 13685–13693 (2006).
91. Swaney, J.S. *et al.* Focal adhesions in (myo)fibroblasts scaffold adenyl cyclase with phosphorylated caveolin. *J. Biol. Chem.* **281**, 17173–17179 (2006).
92. Cheng, X. & Jaggard, J.H. Genetic ablation of caveolin-1 modifies Ca²⁺ spark coupling in vascular arterial smooth muscle cells. *Am. J. Physiol. Heart Circ. Physiol.* **290**, H2309–H2319 (2006).
93. Wang, X.L. *et al.* Caveolae targeting and regulation of large conductance Ca²⁺-activated K⁺ channels in vascular endothelial cells. *J. Biol. Chem.* **280**, 11656–11664 (2005).
94. Trigatti, B.L., Anderson, R.G. & Gerber, G.E. Identification of caveolin-1 as a fatty acid binding protein. *Biochem. Biophys. Res. Commun.* **255**, 34–39 (1999).
95. Martin, S. & Parton, R.G. Lipid droplets: a unified view of a dynamic organelle. *Nature Rev. Mol. Cell Biol.* **7**, 373–378 (2006).
96. Murphy, D.J. The biogenesis and functions of lipid bodies in animals, plants and microorganisms. *Prog. Lipid Res.* **40**, 325–438 (2001).
97. Martin, S. & Parton, R.G. Caveolin, cholesterol, and lipid bodies. *Semin. Cell Dev. Biol.* **16**, 163–174 (2005).
98. Brasaemle, D.L., Dolios, G., Shapiro, L. & Wang, R. Proteomic analysis of proteins associated with lipid droplets of basal and lipolytically stimulated 3T3-L1 adipocytes. *J. Biol. Chem.* **279**, 46835–46842 (2004).
99. Pol, A. *et al.* Dynamic and regulated association of caveolin with lipid bodies: modulation of lipid body motility and function by a dominant negative mutant. *Mol. Biol. Cell* **15**, 99–110 (2004).
100. Meshulam, T., Simard, J.R., Wharton, J., Hamilton, J.A. & Pilch, P.F. Role of caveolin-1 and cholesterol in transmembrane fatty acid movement. *Biochemistry* **45**, 2882–2893 (2006).
101. Fu, Y. *et al.* Expression of caveolin-1 enhances cholesterol efflux in hepatic cells. *J. Biol. Chem.* **279**, 14140–14146 (2004).
102. Fielding, C.J. & Fielding, P.E. Caveolae and intracellular trafficking of cholesterol. *Adv. Drug Deliv. Rev.* **49**, 251–264 (2001).
103. Frank, P.G. *et al.* Caveolin-1 and the regulation of cellular cholesterol homeostasis. *Am. J. Physiol. Heart Circ. Physiol.* **291**, H677–H686 (2006).
104. Pol, A. *et al.* A caveolin dominant negative mutant associates with lipid bodies and induces intracellular cholesterol imbalance. *J. Cell Biol.* **152**, 1057–1070 (2001).
105. Fernandez, M.A. *et al.* Caveolin-1 is essential for liver regeneration. *Science* **313**, 1628–1632 (2006).
106. Razani, B. *et al.* Caveolin-1-deficient mice are lean, resistant to diet-induced obesity, and show hypertriglyceridemia with adipocyte abnormalities. *J. Biol. Chem.* **277**, 8635–8647 (2002).
107. Le Lay, S. *et al.* Cholesterol-induced caveolin targeting to lipid droplets in adipocytes: a role for caveolar endocytosis. *Traffic* **7**, 549–561 (2006).
108. Landth, T. From entangled membranes to eclectic morphologies: cubic membranes as subcellular space organizers. *FEBS Lett.* **369**, 13–17 (1995).
109. Vogel, V. & Sheetz, M. Local force and geometry sensing regulate cell functions. *Nature Rev. Mol. Cell Biol.* **7**, 265–275 (2006).
110. Rizzo, V., Morton, C., DePaola, N., Schnitzer, J.E. & Davies, P.F. Recruitment of endothelial caveolae into mechanotransduction pathways by flow conditioning *in vitro*. *Am. J. Physiol. Heart Circ. Physiol.* **285**, H1720–H1729 (2003).
- In cultured endothelial cells, chronic exposure to shear stress causes increased surface localization of CAV1 and caveolae; this effect is linked to increased mechanosensitivity and activation of specific signalling pathways.**
111. Boyd, N.L. *et al.* Chronic shear induces caveolae formation and alters ERK and Akt responses in endothelial cells. *Am. J. Physiol. Heart Circ. Physiol.* **285**, H1113–H1122 (2003).
112. Radel, C. & Rizzo, V. Integrin mechanotransduction stimulates caveolin-1 phosphorylation and recruitment of Csk to mediate actin reorganization. *Am. J. Physiol. Heart Circ. Physiol.* **288**, H936–H945 (2005).
Cultured bovine aortic endothelial cells subjected to shear stress show rapid Tyr14 phosphorylation of CAV1, which is dependent on β 1-integrin activation, supporting a role for caveolae in mechanosensing.
113. Yu, J. *et al.* Direct evidence for the role of caveolin-1 and caveolae in mechanotransduction and remodeling of blood vessels. *J. Clin. Invest.* **116**, 1284–1291 (2006).
Endothelial expression of CAV1 in Cav1-null mice shows that CAV1 is required for eNOS activation in response to endothelial flow; these data provide further support for a mechanosensing role of caveolae.
114. Rudic, R.D. *et al.* Direct evidence for the importance of endothelium-derived nitric oxide in vascular remodeling. *J. Clin. Invest.* **101**, 731–736 (1998).
115. Sedding, D.G. *et al.* Caveolin-1 facilitates mechanosensitive protein kinase B (Akt) signaling *in vitro* and *in vivo*. *Circ. Res.* **96**, 635–642 (2005).
Smooth-muscle cells subjected to cyclic stretch show rapid redistribution of CAV1 to focal contacts. CAV1 is required for stretch-triggered cell-cycle progression.
116. Oh, P. & Schnitzer, J.E. Segregation of heterotrimeric G proteins in cell surface microdomains. G_i binds caveolin to concentrate in caveolae, whereas G_s and G_q target lipid rafts by default. *Mol. Biol. Cell* **12**, 685–698 (2001).
117. Czarny, M. & Schnitzer, J.E. Neutral sphingomyelinase inhibitor scyphostatin prevents and ceramide mimics mechanotransduction in vascular endothelium. *Am. J. Physiol. Heart Circ. Physiol.* **287**, H1344–H1352 (2004).
118. Aleghat, F.J. & Ingber, D.E. Mechanotransduction: all signals point to cytoskeleton, matrix, and integrins. *Sci. STKE* **2002**, PE6 (2002).
119. Lundbaek, J.A. *et al.* Regulation of sodium channel function by bilayer elasticity: the importance of hydrophobic coupling. Effects of micelle-forming amphiphiles and cholesterol. *J. Gen. Physiol.* **123**, 599–621 (2004).
120. Lundbaek, J.A., Andersen, O.S., Werge, T. & Nielsen, C. Cholesterol-induced protein sorting: an analysis of energetic feasibility. *Biophys. J.* **84**, 2080–2089 (2003).
121. Sens, P. & Turner, M.S. Budded membrane microdomains as tension regulators. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **73**, 031918 (2006).
122. Koleske, A.J., Baltimore, D. & Lisanti, M.P. Reduction of caveolin and caveolae in oncogenically transformed cells. *Proc. Natl Acad. Sci. USA* **92**, 1381–1385 (1995).
123. Lee, S.W., Reimer, C.L., Oh, P., Campbell, D.B. & Schnitzer, J.E. Tumor cell growth inhibition by caveolin re-expression in human breast cancer cells. *Oncogene* **16**, 1391–1397 (1998).
124. Hayashi, K. *et al.* Invasion activating caveolin-1 mutation in human scirrhous breast cancers. *Cancer Res.* **61**, 2361–2364 (2001).
125. Williams, T.M. *et al.* Loss of caveolin-1 gene expression accelerates the development of dysplastic mammary lesions in tumor-prone transgenic mice. *Mol. Biol. Cell* **14**, 1027–1042 (2003).
126. Capozza, F. *et al.* Absence of caveolin-1 sensitizes mouse skin to carcinogen-induced epidermal hyperplasia and tumor formation. *Am. J. Pathol.* **162**, 2029–2039 (2003).
127. Li, T. *et al.* Caveolin-1 mutations in human breast cancer: functional association with estrogen receptor α -positive status. *Am. J. Pathol.* **168**, 1998–2013 (2006).
128. Sunaga, N. *et al.* Different roles for caveolin-1 in the development of non-small cell lung cancer versus small cell lung cancer. *Cancer Res.* **64**, 4277–4285 (2004).
129. Thompson, T.C., Timme, T.L., Li, L. & Goltsov, A. Caveolin-1, a metastasis-related gene that promotes cell survival in prostate cancer. *Apoptosis* **4**, 233–237 (1999).
130. Yang, G., Timme, T.L., Frolov, A., Wheeler, T.M. & Thompson, T.C. Combined c-Myc and caveolin-1 expression in human prostate carcinoma predicts prostate carcinoma progression. *Cancer* **103**, 1186–1194 (2005).
131. Vorgerd, M. *et al.* A sporadic case of rippling muscle disease caused by a *de novo* caveolin-3 mutation. *Neurology* **57**, 2273–2277 (2001).
132. McNally, E.M. *et al.* Caveolin-3 in muscular dystrophy. *Hum. Mol. Genet.* **7**, 871–877 (1998).
133. Woodman, S.E., Sotgia, F., Galbiati, F., Minetti, C. & Lisanti, M.P. Caveolinopathies: mutations in caveolin-3 cause four distinct autosomal dominant muscle diseases. *Neurology* **62**, 538–543 (2004).
134. Minetti, C. *et al.* Mutations in the caveolin-3 gene cause autosomal dominant limb-girdle muscular dystrophy. *Nature Genet.* **18**, 365–368 (1998).
135. Galbiati, F., Volante, D., Minetti, C., Chu, J.B. & Lisanti, M.P. Phenotypic behavior of caveolin-3 mutations that cause autosomal dominant limb girdle muscular dystrophy (LGMD-1C). Retention of LGMD-1C caveolin-3 mutants within the golgi complex. *J. Biol. Chem.* **274**, 25632–25641 (1999).
136. Matsuda, C. *et al.* The sarcolemmal proteins dysferlin and caveolin-3 interact in skeletal muscle. *Hum. Mol. Genet.* **10**, 1761–1766 (2001).
137. Parton, R.G., Molero, J.C., Floetenmeyer, M., Green, K.M. & James, D.E. Characterization of a distinct plasma membrane macrodomain in differentiated adipocytes. *J. Biol. Chem.* **277**, 46769–46778 (2002).
138. Foster, L.J., De Hoog, C.L. & Mann, M. Unbiased quantitative proteomics of lipid rafts reveals high specificity for signaling factors. *Proc. Natl Acad. Sci. USA* **100**, 5813–5818 (2003).
139. Nixon, S.J. *et al.* Zebrafish as a model for caveolin-associated muscle disease; caveolin-3 is required for myofibril organization and muscle cell patterning. *Hum. Mol. Genet.* **14**, 1727–1743 (2005).
140. Ohsawa, Y. *et al.* Overexpression of P104L mutant caveolin-3 in mice develops hypertrophic cardiomyopathy with enhanced contractility in association with increased endothelial nitric oxide synthase activity. *Hum. Mol. Genet.* **13**, 151–157 (2004).
141. Sharma, D.K. *et al.* Glycosphingolipids internalized via caveolar-related endocytosis rapidly merge with the clathrin pathway in early endosomes and form microdomains for recycling. *J. Biol. Chem.* **278**, 7564–7572 (2003).

Acknowledgments

We are extremely grateful to R. Stan, B. Sessa, J. Hancock, and members of the Parton and Simons laboratories for comments on the manuscript. The authors acknowledge the continuing support of the National Health and Medical Research Council of Australia (R.G.P.), the National Institutes of Health, USA (R.G.P.), the Max Planck Society (K.S.), Deutsche Forschungsgemeinschaft (K.S.) and the European Commission (K.S.). The Institute for Molecular Bioscience is a Special Research Centre of the Australian Research Council. We would also like to apologize to those researchers whose primary research could not be cited owing to space constraints.

Competing interests statement

The authors declare no competing financial interests.

DATABASES

The following terms in this article are linked online to:
Entrez gene: <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>
FimH
UniProtKB: <http://us.expasy.org/sprot/CAV1|CAV2|CAV3|dynam|dysferlin|eNOS|fibronectin|ICAM1|RAB5|Rac1|syntaxin-6|TRPC1>

FURTHER INFORMATION

Rob Parton's laboratory homepage:
<http://www.imb.uq.edu.au/index.html?id=11688>
The Simons group laboratory homepage:
<http://www.mpi-cbg.de/research/groups/simons/simons.html>
Access to this links box is available online.